



order conditions (e.g., the concentration of amines was at least 20 times greater than that of the substrate **2**). The reactions were followed by monitoring the appearance of 3,4-dinitrophenoxide at 410 nm. All reactions obeyed first-order kinetics. Pseudo-first-order rate constants ( $k_{\text{obsd}}$ ) were determined from the equation,  $\ln(A - A_T) = -k_{\text{obsd}}t + C$ . The plots of  $k_{\text{obsd}}$  vs. the amine concentration were linear passing through the origin, indicating that general base catalysis by a second amine molecule is absent and the contribution of  $\text{H}_2\text{O}$  and/or  $\text{OH}^-$  ion from the hydrolysis of amines to  $k_{\text{obsd}}$  is negligible. Thus, the rate equation can be given as eq. (1). The apparent second-order rate constants ( $k_N$ ) were determined from the slope of the linear plots of  $k_{\text{obsd}}$  vs. the amine concentration. Generally five different amine concentrations were used to determine  $k_N$  values. It is estimated from replicate runs that the uncertainty in the rate constants is less than 3%. The  $k_N$  values determined in this way are summarized in Table 1 together with those reported for the corresponding reactions of **1** for comparison purpose.

$$\text{rate} = k_{\text{obsd}}[\mathbf{2}], \text{ where } k_{\text{obsd}} = k_N[\text{amine}] \quad (1)$$

**Effect of *ortho*-Nitro Group on Reactivity and Mechanism.** As shown in Table 1, the second-order rate constant for the reaction of **2** decreases as the basicity of amines decreases, i.e.,  $k_N$  decreases from  $152 \text{ M}^{-1}\text{s}^{-1}$  to  $10.1$  and  $0.080 \text{ M}^{-1}\text{s}^{-1}$  as the  $\text{p}K_a$  of amines decreases from 11.02 to 8.65 and 5.95, respectively. A similar result is shown for the corresponding reactions of **1**. Interestingly, **2** is less reactive than **1** when the attacking amines are weakly basic ( $\text{p}K_a < 10.4$ ) but becomes more reactive as the amine basicity increases further.


The effect of amine basicity on reactivity is illustrated in Figure 1. The Brønsted-type plots are curved downwardly for the reactions of **2**, when  $k_N$  and  $\text{p}K_a$  are statistically corrected using  $p$  and  $q$  (i.e.,  $p = 2$  except  $p = 4$  for piperazinium ion and  $q = 1$  except  $q = 2$  for piperazine).<sup>11</sup> A similar result is demonstrated for the corresponding reactions of **1**, although the slopes are slightly different, i.e., the slope of the Brønsted-type plots decreases from 0.83 to 0.34 and from 0.74 to 0.34 for the reactions of **2** and **1**, respectively, as the amine basicity increases.

It has generally been understood that a change in the RDS

**Table 1.** Summary of Second-order Rate Constants ( $k_N$ ,  $\text{M}^{-1}\text{s}^{-1}$ ) for the Reactions of 3,4-Dinitrophenyl 2-Thiophenecarboxylates (**2**) and 2,4-Dinitrophenyl 2-Thiophenecarboxylates (**1**) with Secondary Alicyclic Amines in 20 mol % DMSO at  $25.0 \pm 0.1 \text{ }^\circ\text{C}$ <sup>a</sup>

Entry	Amines	$\text{p}K_a$	$k_N/\text{M}^{-1}\text{s}^{-1}$	
			<b>2</b>	<b>1</b>
1	piperidine	11.02	152	145
2	3-methylpiperidine	10.80	162	139
3	piperazine	9.85	64.1	68.2
4	morpholine	8.65	10.1	15.3
5	1-formylpiperazine	7.98	1.76	4.04
6	piperazinium ion	5.95	0.080	0.397


<sup>a</sup>The data for the reactions of **1** were taken from ref. 10.




**Figure 1.** Brønsted-type plots for the reactions of 2,4-dinitrophenyl 2-thiophenecarboxylates **1** (○) and 3,4-dinitrophenyl 2-thiophenecarboxylates **2** (●) with secondary alicyclic amines in 80 mol %  $\text{H}_2\text{O}/20 \text{ mol } \%$  DMSO at  $25.0 \pm 0.1 \text{ }^\circ\text{C}$ . The identity of points is given in Table 1.

occurs as the attacking amine becomes more basic than the leaving group by 4 to 5  $\text{p}K_a$  units.<sup>1-5</sup> The  $\text{p}K_a$  at the center of Brønsted curvature has been defined as  $\text{p}K_a^0$ .<sup>1d,12</sup> The  $\text{p}K_a^0$  value for the reactions of **1** has been reported to be 9.1, which is ca. 5  $\text{p}K_a$  units larger than the  $\text{p}K_a$  of the conjugate acid of the leaving 2,4-dinitrophenoxide ( $\text{p}K_a = 4.11$ ). Thus, the curved Brønsted-type plot obtained for the aminolysis of **1** has been taken as evidence for a change in RDS of a stepwise reaction.<sup>10</sup> As shown in Figure 1, the  $\text{p}K_a^0$  value for the reactions of **2** is 10.4, which is ca. 5  $\text{p}K_a$  units larger than the  $\text{p}K_a$  of the conjugate acid of the leaving 3,4-dinitrophenoxide ( $\text{p}K_a = 5.42$ ). Accordingly, one can suggest that the current aminolysis of **2** also proceeds through  $\text{T}^\ddagger$  as shown in Scheme 1 and the RDS changes at  $\text{p}K_a = 10.4$ .

To examine the above argument a plot of  $\log k_N$  for the reactions of **1** vs.  $\log k_N$  for the corresponding reactions of **2** has been constructed. As shown in Figure 2, an excellent linear correlation is obtained (e.g.,  $R^2 = 0.9992$ ) with a slope of 1.29. Such a good linear plot suggests that the reactions of **1** and **2** proceed through the same mechanism. The slope of 1.29 is consistent with the fact that the reactions of **2** exhibit



**Scheme 1**



**Figure 2.** Plot of  $\log k_N$  for the reactions of 2,4-dinitrophenyl 2-thiophenecarboxylates (**1**) vs.  $\log k_N$  for the reactions of 3,4-dinitrophenyl 2-thiophenecarboxylates (**2**) in 80 mol % H<sub>2</sub>O/20 mol % DMSO at 25.0 ± 0.1 °C. The identity of numbers is given in Table 1.

a larger slope than those of **1** in the Brønsted-type plots (see Figure 1). Accordingly, one can suggest that shifting the NO<sub>2</sub> group from the *ortho*-position to the *meta*-position of the leaving group does not alter the reaction mechanism but influences the reactivity and the Brønsted coefficient.

**Determination of Microscopic Rate Constants.** On the basis of the mechanism proposed above, the curved Brønsted-type plot for the aminolysis of **2** has been analyzed using a semiempirical equation (eq. 2).<sup>1d,12</sup> The parameters  $\beta_1$  and  $\beta_2$  represent the slope of the curved Brønsted plot for the reactions with strongly basic and weakly basic amines, respectively. Here  $k_N^\circ$  refers to the  $k_N$  value at  $pK_a^\circ$ . The parameters determined from the fitting of eq. (2) to the experimental points are  $\beta_1 = 0.34$ ,  $\beta_2 = 0.83$ , and  $pK_a^\circ = 10.4$  for the reactions of **2**. The  $\beta_1$  value for the reactions of **2** is the same as that reported for the reactions of **1**, while  $\beta_2$  is slightly larger for the reactions of **2** ( $\beta_2 = 0.83$ ) than for those of **1** ( $\beta_2 = 0.74$ ).

$$\log(k_N/k_N^\circ) = \beta_2(pK_a - pK_a^\circ) - \log(1 + \alpha)/2$$

where  $\log \alpha = (\beta_2 - \beta_1)(pK_a - pK_a^\circ)$  (2)

The  $k_N$  values for the reactions of **2** have been dissected into their microscopic rate constants through eqs. (3)-(10) as shown below. The apparent second-order rate constant  $k_N$  can be expressed as eq. (3) by applying the steady-state conditions to the intermediate on the basis of the proposed mechanism.

$$k_N = k_1 k_2 / (k_{-1} + k_2) \quad (3)$$

The  $k_2/k_{-1}$  ratios associated with the aminolysis of **2** have been determined using eqs. (4)-(9). Eq. (3) can be simplified to eq. (4) or (5). Then,  $\beta_1$  and  $\beta_2$  can be expressed as eqs. (6)

**Table 2.** Summary of Microscopic Rate Constants ( $k_1$  and  $k_2/k_{-1}$  Ratios) for the Reactions of 2,4-Dinitrophenyl 2-Thiophenecarboxylates (**1**) and 3,4-Dinitrophenyl 2-Thiophenecarboxylates (**2**) with Alicyclic Secondary Amines in 80 mol % H<sub>2</sub>O/20 mol % DMSO at 25.0 ± 0.1 °C<sup>a</sup>

Entry	Amines	pK <sub>a</sub>	10 <sup>2</sup> k <sub>2</sub> /k <sub>-1</sub>		k <sub>1</sub> /M <sup>-1</sup> s <sup>-1</sup>	
			<b>1</b>	<b>2</b>	<b>1</b>	<b>2</b>
1	piperidine	11.02	773	282	164	206
2	3-methylpiperidine	10.80	631	220	161	236
3	piperazine	9.85	200	53.8	102	183
4	morpholine	8.65	87.1	19.5	32.9	62.0
5	1-formylpiperazine	7.98	47.0	9.15	12.6	21.0
6	piperazinium ion	5.95	9.55	1.30	4.55	6.24

<sup>a</sup>The data for the reactions of **1** were taken from ref. 10.

and (7), respectively.

$$k_N = k_1 k_2 / k_{-1}, \text{ when } k_2 \ll k_{-1} \quad (4)$$

$$k_N = k_1, \text{ when } k_2 \gg k_{-1} \quad (5)$$

$$\beta_1 = d(\log k_1) / d(pK_a) \quad (6)$$

$$\beta_2 = d(\log k_1 k_2 / k_{-1}) / d(pK_a)$$

$$= \beta_1 + d(\log k_2 / k_{-1}) / d(pK_a) \quad (7)$$

Eq. (7) can be rearranged as eq. (8). Integral of eq. (8) from  $pK_a^\circ$  results in eq. (9). Since  $k_2 = k_{-1}$  at  $pK_a^\circ$ , the term  $(\log k_2 / k_{-1})_{pK_a^\circ}$  is zero. Therefore, one can calculate the  $k_2/k_{-1}$  ratios for the aminolysis of **2** from eq. (9) using  $pK_a^\circ = 10.4$ ,  $\beta_1 = 0.34$ , and  $\beta_2 = 0.83$ . The  $k_1$  values have been determined from eq. (10) using the  $k_N$  values in Table 1 and the  $k_2/k_{-1}$  ratios calculated above. The  $k_2/k_{-1}$  ratios and  $k_1$  values are summarized in Table 2.

$$\beta_2 - \beta_1 = d(\log k_2 / k_{-1}) / d(pK_a) \quad (8)$$


$$(\log k_2 / k_{-1})_{pK_a} = (\beta_2 - \beta_1)(pK_a - pK_a^\circ) \quad (9)$$

$$k_N = k_1 k_2 / (k_{-1} + k_2) = k_1 / (k_{-1} / k_2 + 1) \quad (10)$$

**Effect of *ortho*-Nitro Group on Microscopic Rate Constants.** As shown in Table 2, the  $k_2/k_{-1}$  ratio decreases as the amine basicity decreases for both reactions of **1** and **2**. The  $k_2$  value has been suggested to be independent of the basicity of amines, since the N atom of the aminium moiety of T<sup>±</sup> cannot provide a push to expel the leaving aryloxide from T<sup>±</sup> due to the lack of an electron pair on its nitrogen atom.<sup>1d,13</sup> On the other hand, the  $k_{-1}$  value would increase as the amine basicity decreases. Thus, one can expect that the  $k_2/k_{-1}$  ratio would decrease as the amine basicity decreases. In fact, as shown in Table 2, the  $k_2/k_{-1}$  ratio decreases as the amine basicity decreases.


In Figure 3 is illustrated the effect of amine basicity on the  $k_2/k_{-1}$  ratio. The plots of  $\log k_2/k_{-1}$  vs.  $pK_a$  of the conjugate acid of amines are linear for the reactions of **1** and **2**, although the slope of the linear plots is slightly larger for the reactions of **2** (*i.e.*,  $\beta_1 = 0.49$ ) than for those of **1** (*i.e.*,  $\beta_1 = 0.40$ ).

The reaction of **2** would result in a smaller  $k_2$  value than



**Figure 3.** Plots of  $\log k_2/k_{-1}$  versus  $pK_a$  for the reactions of 2,4-dinitrophenyl 2-thiophenecarboxylates **1** ( $\circ$ ) and 3,4-dinitrophenyl 2-thiophenecarboxylates **2** ( $\bullet$ ) with secondary alicyclic amines in 80 mol %  $H_2O$ /20 mol % DMSO at  $25.0 \pm 0.1$  °C. The identity of points is given in Table 2.

that of **1** for a given amine, since 3,4-dinitrophenoxide is more basic and a poorer nucleofuge than 2,4-dinitrophenoxide. On the other hand, the  $k_{-1}$  value would be little influenced by the basicity of the leaving group. Thus, one might expect that the  $k_2/k_{-1}$  ratio is smaller for the reaction of **2** than for the corresponding reaction of **1**. In fact, Table 2



**Figure 4.** Brønsted-type plots for the reactions of 2,4-dinitrophenyl 2-thiophenecarboxylates **1** ( $\circ$ ) and 3,4-dinitrophenyl 2-thiophenecarboxylates **2** ( $\bullet$ ) with secondary alicyclic amines in 80 mol %  $H_2O$ /20 mol % DMSO at  $25.0 \pm 0.1$  °C. The identity of points is given in Table 2.

and Figure 3 show that the  $k_2/k_{-1}$  ratio is smaller for the reaction of **2** than for the reaction of **1** for a given amine.

Table 2 shows that  $k_1$  increases with increasing amine basicity for the reactions of **1** and **2**. The effect of amine basicity on  $k_1$  is illustrated in Figure 4. The plots of  $\log k_1$  vs.  $pK_a$  of the conjugate acid of amines are linear. Interestingly, the reaction of **2** exhibits a slightly larger  $k_1$  value than that of **1** for a given amine. This result is unexpected, since the C=O bond of **2** would be less electrophilic than that of **1** on the basis of the fact that 3,4-dinitrophenoxide is more basic than 2,4-dinitrophenoxide. Accordingly, the difference in basicity between the two nucleofuges cannot account for the  $k_1$  values determined for the reactions of **1** and **2**.

The steric hindrance exerted by the *ortho*- $NO_2$  in substrate **1** would be significant for the nucleophilic attack process (or when the  $k_1$  step is the RDS). This idea is consistent with the fact that  $k_1$  is smaller for the reactions of **1** than for the corresponding reactions of **2**, and the reactions of **1** with strongly basic piperidine and 3-methylpiperidine result in smaller  $k_N$  values than for the corresponding reactions of **2** when the  $k_1$  step is the RDS. However, the steric hindrance appears to be insignificant when the  $k_2$  step is the RDS, since the reactions of **1** with weakly basic amines result in larger  $k_N$  values than for the corresponding reactions of **2** (note that the  $k_2$  step is the RDS when  $pK_a < 10.4$ ). Thus, one can suggest that steric hindrance is mainly responsible for the fact that the reaction of **1** results in a smaller  $k_N$  value than the corresponding reaction of **2** when  $pK_a > 10.4$  (or  $k_1$  is smaller for the former reactions than for the latter ones regardless of the amine basicity).

## Conclusions

The current study has allowed us to conclude the following: (1) Aminolyses of **1** and **2** proceed through  $T^\ddagger$  with a change in the RDS. (2) Substrate **2** is less reactive than substrate **1** toward weakly basic amines ( $pK_a < 10.4$ ) but becomes more reactive as the basicity of amines increases further. (3) Dissection of  $k_N$  into the microscopic rate constants has revealed that aminolysis of **2** results in smaller  $k_2/k_{-1}$  ratio as expected, however unexpectedly, the  $k_1$  value is larger for the reaction of **2** than the corresponding reaction of **1**. (4) The steric effect of the *ortho*- $NO_2$  is significant when the  $k_1$  is the RDS but insignificant when the  $k_2$  process is the slow step. (5) The steric hindrance exerted by the *ortho*- $NO_2$  is mainly responsible for the fact that the reaction of **1** results in a smaller  $k_N$  value than the corresponding reaction of **2** when  $pK_a > 10.4$  (or  $k_1$  is smaller for the former reactions than for the latter ones regardless of the amine basicity).

## Experimental Section

**Materials.** Substrate **2** was readily prepared from the reaction of 3,4-dinitrophenol and 2-thiophenecarbonyl chloride in the presence of triethylamine in anhydrous ether. The purity of **2** was confirmed by its melting point and  $^1H$

NMR spectrum. Due to the low solubility of **2** in pure H<sub>2</sub>O, 20 mol % DMSO/80 mol % H<sub>2</sub>O was used as the reaction medium. Doubly glass distilled water was further boiled and cooled under nitrogen just before use.

**Kinetics.** The kinetic studies were performed at 25.0 ± 0.1 °C with a Scinco S-3100 UV-Vis spectrophotometer equipped with a constant temperature circulating bath for slow reactions (*e.g.*,  $t_{1/2} \geq 10$  s) or with an Applied Photophysics Stopped-flow spectrophotometer for fast reactions (*e.g.*,  $t_{1/2} < 10$  s). The reactions were followed by monitoring the appearance of 3,4-dinitrophenoxide ion at 410 nm. All the reactions were carried out under pseudo-first-order conditions in which the amine concentration was at least 20 times greater than that of the substrate **2**.

**Products Analysis.** 3,4-Dinitrophenoxide ion was liberated quantitatively and identified as one of the reaction products by comparison of the UV-vis spectra after the completion of the reactions with those of the authentic samples under the same reaction conditions.

**Acknowledgments.** This work was supported by a grant from Korea Research Foundation (KRF-2005-015-C00256). Jin-A Seo is also grateful for the BK 21 scholarship.

## References

- (a) Jencks, W. P. *Catalysis in Chemistry and Enzymology*; McGraw-Hill: New York, 1969; pp 480-483. (b) Jencks, W. P. *J. Chem. Soc. Rev.* **1981**, *10*, 345-375. (c) Jencks, W. P.; Gilchrist, M. *J. Am. Chem. Soc.* **1968**, *90*, 2622-2637. (d) Gresser, M. J.; Jencks, W. P. *J. Am. Chem. Soc.* **1977**, *99*, 6963-6970.
- (a) Williams, A. *Adv. Phys. Org. Chem.* **1992**, *27*, 2-55. (b) Menger, F. M.; Smith, J. H. *J. Am. Chem. Soc.* **1972**, *94*, 3824-3829. (c) Tsang, W. Y.; Ahmed, N.; Page, M. I. *Org. Biomol. Chem.* **2007**, *5*, 485-493.
- (a) Castro, E. A.; Echevarria, G. R.; Opazo, A.; Robert, P.; Santos, J. G. *J. Phys. Org. Chem.* **2006**, *19*, 129-135. (b) Castro, E. A.; Campodonico, P. R.; Contreras, R.; Fuentealba, P.; Santos, J. G.; Leis, J. R.; Garcia-Rio, L.; Saez, J. A.; Domingo, L. R. *Tetrahedron* **2006**, *62*, 2555-2562. (c) Castro, E. A.; Aliaga, M.; Gazitua, M.; Santos, J. G. *Tetrahedron* **2006**, *62*, 4863-4869. (d) Castro, E. A.; Aguayo, R.; Bessolo, J.; Santos, J. G. *J. Org. Chem.* **2005**, *70*, 7788-7791. (e) Castro, E. A.; Aliaga, M.; Santos, J. G. *J. Org. Chem.* **2005**, *70*, 2679-2685. (f) Castro, E. A.; Gazitua, M.; Santos, J. G. *J. Org. Chem.* **2005**, *70*, 8088-8092.
- (a) Sung, D. D.; Kang, S. S.; Lee, J. P.; Jung, D. I.; Ryu, Z. H.; Lee, I. *Bull. Korean Chem. Soc.* **2007**, *28*, 1670-1674. (b) Hoque, M. E. U.; Dey, N. K.; Guha, A. K.; Kim, C. K.; Lee, B. S.; Lee, H. W. *Bull. Korean Chem. Soc.* **2007**, *28*, 1797-1802. (c) Kim, C. K.; Kim, D. J.; Zhang, H.; Hsieh, Y. H.; Lee, B. S.; Lee, H. W.; Kim, C. K. *Bull. Korean Chem. Soc.* **2007**, *28*, 1031-1034. (d) Ehtesham, M. H. U.; Lee, H. W. *Bull. Korean Chem. Soc.* **2007**, *28*, 936-940. (e) Sung, D. D.; Koo, I. S.; Yang, K. Y.; Lee, I. *Chem. Phys. Lett.* **2006**, *432*, 426-430. (f) Sung, D. D.; Koo, I. S.; Yang, K. Y.; Lee, I. *Chem. Phys. Lett.* **2006**, *426*, 280-284.
- (a) Um, I. H.; Lee, J. Y.; Ko, S. H.; Bae, S. K. *J. Org. Chem.* **2006**, *71*, 5800-5803. (b) Um, I. H.; Lee, J. Y.; Lee, H. W.; Nagano, Y.; Fujio, M.; Tsuno, Y. *J. Org. Chem.* **2005**, *70*, 4980-4987. (c) Um, I. H.; Kim, K. H.; Park, H. R.; Fujio, M.; Tsuno, Y. *J. Org. Chem.* **2004**, *69*, 3937-3942.
- Um, I. H.; Jeon, S. E.; Seok, J. A. *Chem. Eur. J.* **2006**, *12*, 1237-1243.
- (a) Um, I. H.; Park, Y. M.; Fujio, M.; Mishima, M.; Tsuno, Y. *J. Org. Chem.* **2007**, *72*, 4816-4821. (b) Um, I. H.; Akhtar, K.; Park, Y. M.; Khan, S. *Bull. Korean Chem. Soc.* **2007**, *28*, 1353-1357.
- (a) *Techniques of Organic Chemistry*, 4rd ed.; Bernasconi, C. F., Ed.; Wiley: New York, 1986; vol. 6. (b) *Techniques of Organic Chemistry*, 3rd ed.; Lewis, E. S., Ed.; Wiley: New York, 1974; Vol. 6, Part 1. (c) *Advances in Linear Free Energy Relationship*; Chapman, N. B.; Shorter, J., Eds.; Plenum: London, 1972. (d) Pross, A. *Advances in Physical Organic Chemistry*; Academic Press: London, 1977; Vol. 14, pp 69-132.
- (a) Um, I. H.; Shin, Y. H.; Han, J. Y.; Mishima, M. *J. Org. Chem.* **2006**, *71*, 7715-7720. (b) Um, I. H.; Akhtar, K.; Shin, Y. H.; Han, J. Y. *J. Org. Chem.* **2007**, *72*, 3823-3829.
- Um, I. H.; Min, S. W.; Chun, S. M. *Bull. Korean Chem. Soc.* **2008**, submitted for publication.
- Bell, R. P. *The Proton in Chemistry*; Methuen: London, 1959; p 159.
- Castro, E. A.; Moodie, R. B. *J. Chem. Soc., Chem. Commun.* **1973**, 828-829.
- (a) Castro, E. A.; Santander, C. L. *J. Org. Chem.* **1985**, *50*, 3595-3600. (b) Castro, E. A.; Valdivia, J. L. *J. Org. Chem.* **1986**, *51*, 1668-1672. (c) Castro, E. A.; Steinfert, G. B. *J. Chem. Soc., Perkin Trans. 2* **1983**, 453-457.