

Oxidation of Benzylic Alcohols with Iodine and Lithium Carbonate in Ionic Liquid

Seung Bae Lee and Jong Chan Lee*

Department of Chemistry, Chung-Ang University, Seoul 156-756, Korea. *E-mail: jcleee@cau.ac.kr

Received August 6, 2009, Accepted October 19, 2009

Key Words: Benzylic alcohol, Iodine, Ionic liquid, Oxidation

The oxidation of benzylic alcohols to the corresponding carbonyl compounds is a very important transformation in organic chemistry and a variety of methods have been reported. Numerous reagents have been utilized for the oxidation of benzylic alcohols. For examples transition metals complexes such as chromium(VI) complexes,¹ ruthenium reagents,² and manganese dioxide³ are widely utilized. In addition, hypervalent iodine(V) compounds such as Dess-Martin periodinate⁴ and *o*-iodoxybenzoic acid,⁵ iodosobenzene,⁶ and (diacetoxyiodo)benzene⁷ received much attention due to their outstanding reactivities. However, above mentioned reagents suffered from several drawbacks in terms of harsh reaction conditions, high cost, instability, high toxicity, and operational difficulty. Molecular iodine can be considered one of the most ideal oxidant in organic reactions due to its many desirable properties like low cost, safe to handle,

and ready availability. However, the utilization of iodine for the oxidation of alcohols to the corresponding carbonyl compounds has been very scarce. Furthermore, due to the intrinsic low reactivity of molecular iodine compared to the other highly reactive hypervalent iodine reagents, oxidation of alcohols with molecular iodine generally required the concomitant use of activating agents. For examples, previously reported iodine induced oxidative methods had been invariably carried out in the presence of activating substances such as 2,2,6,6-tetramethyl-1-piperdinyloxy (TEMPO),⁸ potassium iodide,⁹ or (diacetoxyiodo)benzene.¹⁰ To the best of our knowledge, successful utilization of molecular iodine in the absence of any additional promoters in the oxidation of benzylic alcohols to the corresponding carbonyl compounds has been unprecedented.


Herein, we wish to report our findings on the facile oxidation

Table 1. Oxidation of Alcohols to Aldehydes and Ketones with I₂/Li₂CO₃

Entry	Substrate	Product	Time (h)/Temp (°C)	Yield ^a (%)
1	PhCH ₂ OH	$\text{Ph}\overset{\text{O}}{\parallel}\text{CH}$	6/60	85
2	<i>p</i> -CH ₃ C ₆ H ₄ CH ₂ OH	<i>p</i> -CH ₃ C ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CH}$	4/60	81
3	<i>p</i> -FC ₆ H ₄ CH ₂ OH	<i>p</i> -FC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CH}$	4/60	81
4	<i>p</i> -ClC ₆ H ₄ CH ₂ OH	<i>p</i> -ClC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CH}$	4/60	82
5	<i>p</i> -BrC ₆ H ₄ CH ₂ OH	<i>p</i> -BrC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CH}$	4/60	81
6	<i>p</i> -CH ₃ OC ₆ H ₄ CH ₂ OH	<i>p</i> -CH ₃ OC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CH}$	24/60	83
7	Piperonyl alcohol	Piperonyl aldehyde	24/60	71
8	PhCH(CH ₃)OH	Ph $\overset{\text{O}}{\parallel}\text{CCH}_3$	12/r.t	88
9	PhCH(CH ₂ CH ₃)OH	Ph $\overset{\text{O}}{\parallel}\text{CCH}_2\text{CH}_3$	12/r.t	83
10	<i>p</i> -ClC ₆ H ₄ CH(CH ₃)OH	<i>p</i> -ClC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CCH}_3$	12/r.t	87
11	PhCH(CN)OH	Ph $\overset{\text{O}}{\parallel}\text{CCN}$	12/r.t	81
12	(Ph) ₂ CHOH	(Ph) ₂ C=O	12/r.t	84
13	<i>p</i> -CH ₃ OC ₆ H ₄ CH(CH ₃)OH	<i>p</i> -CH ₃ OC ₆ H ₄ $\overset{\text{O}}{\parallel}\text{CCH}_3$	12/r.t	81

^aIsolated yields.

of benzylic alcohols using I_2/Li_2CO_3 in ionic liquid as a useful oxidizing system for the oxidation of benzylic alcohols. Thus, reaction of benzyl alcohol and iodine (1.5 equiv) at 60 °C for 6 h in the presence of lithium carbonate (1.0 equiv) in 1-butyl-3-methylimidazolium tetrafluoroborate, [bmim]BF₄, exclusively yielded the benzaldehyde in 85%. Replacement of Li_2CO_3 by other bases such as K_2CO_3 , Et_3N , and 2,6-lutidine gave always inferior yields of desired products. At the present reaction conditions, various primary benzylic alcohols underwent smooth



oxidation to provide high yields of the corresponding aldehydes as summarized in Table 1. Over-oxidized products were not detected at the present reaction conditions. Next, our attention was directed towards oxidation of secondary alcohols. As exemplified in Table 1, secondary benzylic alcohols (entries 8-13) reacted efficiently giving high yields of ketones. Interestingly, in cases of oxidation of secondary benzylic alcohols, strong temperature dependence of selectivity of reactions were found with the highest at room temperature and somewhat lowered yields were obtained at increased temperatures. At the temperature of 60 °C complicated product mixtures were obtained. Electron-donating and withdrawing substituent in aromatic rings had relatively minor influence on the outcome of the oxidation of substituted benzylic alcohols. In all the attempted examples, our conditions were highly selective and efficient, and no over oxidation products were detected. This appears to be the first example of the oxidation of benzylic alcohols as performed by only molecular iodine without any additional activation reagent.

In conclusion, we have demonstrated a simple and effective transition metal-free protocol for the oxidation of benzylic alcohols to the corresponding aldehydes and ketones. The simple procedure together with use of readily available, safe to use, and

inexpensive molecular iodine should make present protocol an important alternative to other classical methods.

Experimental Section

The 1-butyl-3-methylimidazolium tetrafluoroborate, [bmim]BF₄, was obtained from Aldrich. Thin layer chromatography was performed on Merck silica gel 60 F-254 plates (0.25 mm). Merck silica gel 60 (230 - 400 mesh) was used for flash column chromatography.

General procedure. Benzylic alcohol (1.0 mmol) was added to a well-stirred mixture of iodine (0.381 g, 1.5 mmol) and lithium carbonate (0.073 g, 1.0 mmol) in solution of [bmim]BF₄ (1 mL). The mixture was stirred at 60 °C over 6 h. On completion of the reaction, the mixture was dissolved in dichloromethane (40 mL) and washed with 5% sodium thiosulfate (20 mL). The organic layer was separated and dried over $MgSO_4$ and evaporated under reduced pressure. The residue was purified by flash column chromatography (EtOAc : Hexane = 1 : 3) on silica gel to obtain pure carbonyl compounds.

Acknowledgments. This research was supported by the Chung-Ang University Research Grants in 2009.

References

- Hajipour, A. R.; Mallakpour, S. E.; Khoee, S. *Synlett* **2000**, 740.
- Denmark, S. E.; Cramer, C. J.; Sternberg, J. A. *Helv. Chim. Acta* **1986**, *69*, 1971.
- Uchiyama, M.; Kimura, Y.; Ohta, A. *Tetrahedron Lett.* **2000**, *41*, 10013.
- Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277.
- Ozanne, A.; Pouysegue, L.; Deoernet, D.; Francois, B.; Quideau, S. *Org. Lett.* **2003**, *5*, 2903.
- Yakoo, T.; Matsumoto, K.; Oshima, K. *Chem. Lett.* **1993**, 571.
- Narakasa, K.; Morikawa, A.; Siago, K.; Mukuiyama, T. *Bull. Chem. Soc. Jpn.* **1999**, *55*, 2773.
- Miller, R. A.; Hoerner, R. P. *Org. Lett.* **2003**, *5*, 285.
- Gogoi, P.; Konwar, D. *Org. Biomol. Chem.* **2005**, *3*, 3473.
- Karade, N. N.; Tiwari, G. B.; Huple, D. B. *Synlett* **2005**, 2039.