

Mechanistic Studies on the Photohydration of 1-Aryl-5,5-dimethyl-1,3-hexadiynes

Sang Chul Shim*, Yun Sook Chae, and Eun Kyung Back

Department of Chemistry, the Korea Advanced Institute of Science and Technology, 373-1 Kusung-Dong, Yuseong-Gu, Taejeon 305-701, Korea
Received January 27, 1997

We have recently shown that irradiation of 1-aryl-5,5-dimethyl-1,3-hexadiynes substituted by other than nitro group in aqueous sulfuric acid yielded two types of alkynyl and allenyl ketones through both S_1 and T_1 excited states while nitro-substituted diynes gave only allenyl ketones via T_1 excited states (Scheme 1).¹ The diynes substituted by electron-withdrawing group such as *p*-CO₂CH₃, *m*- and *p*-CF₃ gave only type C allenyl ketones while diynes which have electron-donating groups such as *p*-OCH₃, yielded type D allenyl ketones regioselectively.

In this paper we report substituent effects on the photohydration of 1-aryl-5,5-dimethyl-1,3-hexadiynes focusing on the multiplicity of the reactive excited state.

9-Fluorenone-1-carboxylic acid ($E_T=50.0$ kcal mol⁻¹)² efficiently quenched the photohydration of 1- (*p*-methoxy-carbonylphenyl)-5,5-dimethyl-1,3-hexadiyne (*p*-MCPDHD), 1- (*p*-methoxyphenyl)-5,5-dimethyl-1,3-hexadiyne (*p*-MPDHD), 1- (*m*-trifluoromethylphenyl)-5,5-dimethyl-1,3-hexadiyne (*m*-TPDHD), and 1- (*p*-trifluoromethylphenyl)-5,5-dimethyl-1,3-hexadiyne (*p*-TPDHD). The Stern-Volmer plots for the three photohydration products of *p*-MCPDHD (Type A, B and C) are curved reaching a plateau at about $\Phi_{\text{pdt}}^{\circ} / \Phi_{\text{pdt}} = 1.23, 1.48, \text{ and } 1.54$, respectively, supporting the contribution of both the singlet and triplet excited states to the photohydration (Figure 1).³ Similar non-linear Stern-Volmer plots which have different convergence values ($\Phi_{\text{pdt}}^{\circ} / \Phi_{\text{pdt}} = 1.18\text{--}3.5$) were obtained for *p*-MPDHD, *m*-TPDHD, and *p*-TPDHD.

The total, singlet and triplet photohydration quantum yields (Φ° , Φ_s° and Φ_T°) of the diynes were measured⁴ and tabulated in Table 1 with the calculated and measured ratios of $\Phi_T^{\circ} / \Phi_s^{\circ}$. Most of the values of $\Phi_T^{\circ} / \Phi_s^{\circ}$ are less than

unity indicating that contribution of the singlet excited states to photohydration is greater than that of the triplet excited states.





Figure 1. Stern-Volmer plots for the quenching of photohydration reactions of *p*-MCPDHD by 9-fluorenone-1-carboxylic acid.

Table 1. The quantum yields (Φ° , Φ_s° and Φ_T°)^a for the photohydration reaction of 1-aryl-5,5-dimethyl-1,3-hexadiynes and calculated and measured $\Phi_T^{\circ} / \Phi_s^{\circ}$

X	Products ^b	Φ° ($\times 10^{-2}$)	Φ_s° ($\times 10^{-2}$)	Φ_T° ($\times 10^{-2}$)	$\Phi_T^{\circ} / \Phi_s^{\circ}$ (calcd.) ^c	$\Phi_T^{\circ} / \Phi_s^{\circ}$ (measd.)
<i>p</i> -CO ₂ CH ₃	Type A	1.6	1.3	0.30	0.23	0.24
	Type B	1.6	1.1	0.47	0.48	0.44
	Type C	0.31	0.18	0.13	0.54	0.76
<i>p</i> -OCH ₃	Type A	3.7	3.1	0.60	0.18	0.20
	Type B	2.4	1.6	0.75	0.44	0.47
	Type D	4.9	2.6	2.3	0.89	0.89
<i>m</i> -CF ₃	Type A	0.64	0.29	0.35	1.0	1.2
	Type B	5.0	1.4	3.6	2.4	2.5
	Type C	0.13	0.10	0.030	0.28	0.30
<i>p</i> -CF ₃	Type A	0.60	0.29	0.31	0.95	1.1
	Type B	0.50	0.26	0.24	0.90	0.90
	Type C	0.32	0.20	0.12	0.48	0.60

^a Φ_s° was measured from the quantitative analyses by HPLC after irradiation in the presence of a triplet quencher, 9-fluorenone-1-carboxylic acid. Light intensity was measured by ferrioxalate actinometry. $\Phi_T^{\circ} = \Phi^{\circ} - \Phi_s^{\circ}$, where Φ° is total photohydration quantum yield which is obtained after irradiation in the absence of a triplet quencher. ^bSee Scheme 1. ^cCalculated $\Phi_T^{\circ} / \Phi_s^{\circ}$ was from the convergence value of the corresponding Stern-Volmer plots for the quenching of photohydration reactions by 9-fluorenone-1-carboxylic acid.



Scheme 1.

*To whom all correspondence should be addressed.

As reported previously, the preference of protonation site in the photohydration reaction is dependent on the substituent on the aryl ring. Table 2 shows the ratios of total, singlet and triplet quantum yields for the photohydration through protonation at C_4 or C_1 . It is evident that C_4 is protonated predominantly rather than C_1 when X is electron-donating OCH_3 group ($(\Phi^{\circ})_{C_4}/(\Phi^{\circ})_{C_1}=3.6$). On the other hand, the protonation occurs mainly at C_1 when X is electron-withdrawing group ($(\Phi^{\circ})_{C_4}/(\Phi^{\circ})_{C_1}<1$) and the preference of C_1 protonation is greater in *meta*-substituted than in *para*-substituted diynes when X is CF_3 . This is in conform with the "meta-effect".⁵⁻⁷ The electron-withdrawing effect of substituents in the photohydration is in the order of *m*- $CF_3 > p$ - CO_2CH_3 .

The photohydration reactions of conjugated diynes normally proceed through cumulene-type singlet and/or triplet excited state.⁸⁻¹⁰ In 1-aryl-1,3-diynes, charge separation of the lowest singlet excited state seems to be dichotomic (C_1


or C_4 is positively charged) while S_1 state of arylacetylenes has zwitterionic character with positively charged C_1 and negatively charged C_2 .^{11,12} The T_1 state of 1-aryl-1,3-diynes is weakly charged⁹ even though T_1 of acetylenes is normally considered to be similar to that of diradicals. The proposed mechanism for the reactions involves the protonation step being the rate determining step in the formation of alkynyl ketone products and synchronous H_3O^+ addition mechanism for the formation of allenyl ketones.^{9,10}

A possible charge distribution of excited states is shown in Scheme 2. We can propose the A1 and A2 structures for S_1 state and B1 and B2 structures for T_1 state from the fact that four types of products are formed through both S_1 and T_1 excited states which are known to have a cumulene-type structure. Type A and B products (C_4 protonation products) would be formed from A2 and B2 and type B and C products (C_1 protonation products) would be from A1 and B1. The preference of C_1 protonation for electron-withdrawing groups and C_4 protonation for electron-donating groups is due to the degree of stabilization of negative charge on C_1 in the A1 structure. However, the stabilization effect on the diynes is much smaller than that of arylacetylene because of the delocalization of charge through the conjugated system and the regioselective formation of alkynyl ketones is not observed. The exact reasons why the formation of allenyl ketone is regioselective but that of alkynyl ketones is not and why allenyl ketones are only formed in the case of $X=NO_2$ are not clear. However, the results observed evidently show that regioselectivity and multiplicity of the reactive state in the photohydration of 1-aryl-5,5-dimethyl-1,3-hexadiynes are dependent on the substituents on the aryl ring. The substituents such as *p*- CO_2CH_3 , *m*- and *p*- CF_3 , *m*- and *p*- NO_2 groups showed electron-withdrawing effects on the photohydration reaction and the strength of electron-withdrawing effect was *m*- $NO_2 > p$ - $NO_2 > m$ - $CF_3 > p$ - $CF_3 > p$ - CO_2CH_3 .

Table 2. The ratios of total, singlet and triplet quantum yields for the photohydration of 1-aryl-5,5-dimethyl-1,3-hexadiynes proceeding via protonation at C_4 or C_1

X	$(\Phi^{\circ})_{C_4}/(\Phi^{\circ})_{C_1}$	$(\Phi^{\circ}_s)_{C_4}/(\Phi^{\circ}_s)_{C_1}$	$(\Phi^{\circ}_T)_{C_4}/(\Phi^{\circ}_T)_{C_1}$
<i>p</i> - CO_2CH_3	0.84	1.0	0.5
<i>p</i> - OCH_3	3.6	3.6	3.9
<i>m</i> - CF_3	0.12	0.19	0.097
<i>p</i> - CF_3	0.73	0.63	0.86

$^a(\Phi^{\circ})_{C_4}/(\Phi^{\circ})_{C_1} = \Phi^{\circ}(\text{Type A and D}) / \Phi^{\circ}(\text{Type B and C})$ where Φ° is a total photohydration quantum yield shown in Table 1. For each type of photohydration product, see Scheme 1. $^b(\Phi^{\circ}_s)_{C_4}/(\Phi^{\circ}_s)_{C_1} = \Phi^{\circ}_s(\text{Type A and D}) / \Phi^{\circ}_s(\text{Type B and C})$ where Φ°_s is a photohydration quantum yield via singlet excited state. $^c(\Phi^{\circ}_T)_{C_4}/(\Phi^{\circ}_T)_{C_1} = \Phi^{\circ}_T(\text{Type A and D}) / \Phi^{\circ}_T(\text{Type B and C})$ where Φ°_T is a photohydration quantum yield via triplet excited state.



Scheme 2.

while *p*-OCH₃ group showed electron-donating effect. Most of the substituents favored the singlet excited state in the photohydration of diynes but nitro group favored triplet excited state due to the efficient intersystem crossing.¹³

Acknowledgment. This investigation was supported by the Basic Science Research Institute Program, Korean Ministry of Education (Project No. BSRI-96-3406) and the Korea Science and Engineering Foundation (Grant No. 96-0501-09-01-3).

References

1. Shim, S. C.; Chae, Y. S.; Back, E. K.; Park, S. K. *J. Photochem. Photobiol. A: Chem.* Submitted.
2. Scaiano, J. C. *CRC Handbook of Organic Photochemistry*; CRC Press Inc.: Florida, 1989; p 382.
3. Turro, N. J. *Modern Molecular Photochemistry*; Benjamin/Cummings: Menlo Park, CA, 1978; p 252.
4. (a) Parker, C. A. *Proc. Roy. Soc. (London)* **1953**, A220, 104. (b) Hatchard, C. G.; Parker, C. A. *Proc. Roy. Soc. (London)* **1956**, A235, 518.
5. McEwen, J.; Yates, K. *J. Phy. Org. Chem.* **1991**, 4, 193.
6. Zimmerman, H. E.; Somasekhara, S. *J. Am. Chem. Soc.* **1963**, 85, 922.
7. Zimmerman, H.; Sandel, V. R. *J. Am. Chem. Soc.* **1963**, 85, 915.
8. Shim, S. C.; Lee, T. S. *J. Chem. Soc., Perkin Trans. 2*, **1990**, 1739.
9. Back, E. K.; Shim, S. C. *J. Phy. Org. Chem.* **1995**, 8, 699.
10. Back, E. K.; Lee, S. T.; Chae, Y. S.; Shim, S. C. *J. Photosci.* **1995**, 2, 73.
11. Demoulin, D. *J. Chem. Soc.* **1953**, 2288.
12. Wan, P.; Culshaw, S.; Yates, K. *J. Am. Chem. Soc.* **1982**, 104, 2509.
13. (a) Morrison, H. A. *The Chemistry of the Nitro and Nitroso Groups*; Wiley: New York, 1969; p 165. (b) Hurley, R.; Testa, A. C. *J. Am. Chem. Soc.* **1968**, 90, 1949.

Synthesis and Ionophoric Properties of Mono-Penta Type Mixed-Functionalized Ligands Based-upon *p*-tert- Butylcalix[6]arene

Taehoon Kim, Hyun Jae Cho, Weon Seok Oh,
Sangdoon Ahn[†], Jo Woong Lee^{†*}, and Suk-Kyu Chang*

Department of Chemistry, Chung-Ang University, Seoul 156-756, Korea

[†]Department of Chemistry, Seoul National University, Seoul 151-742, Korea

Received January 31, 1997

Ester, amide, and ether functional groups have been ingeniously utilized by nature for the construction of many bioactive molecules having ionophoric properties. Representative of such ingenuity are valinomycin and related antibiotics.¹ There have been many attempts to mimic the ionophoric properties of these natural antibiotics for the development of new host systems.² Calixarenes are one of the most attractive and widely studied compounds for this purpose and many effective ligating groups have been incorporated into their molecular frameworks.³ However, in order to design more versatile calixarene-based host molecules useful for practical purposes, such as preparation of separation media and sensory materials, more elaborate structural transformations are necessary.⁴ In view of this, we prepared a series of mono-penta type mixed ligating ionophores based upon pentaethyl ester of *p*-tert- butylcalix[6] arene and investigated their ionophoric behaviors toward alkali metal cations.

Mono-penta type mixed ligands were prepared by selective monoalkylation for the introduction of mono-part substituent followed by exhaustive alkylation with ethyl bromoacetate (Scheme). Monoalkylation was performed by reacting with one or two equivalents of required methyl bromoacetate, chloroacetone, or *N,N*- diethyl bromoacetamide in

the presence of K₂CO₃ in THF according to the reported procedure to obtain monomethyl or monobenzyl ether of *p*-tert- butylcalix[6]arene (yield: 42-58%).⁵ The desired mixed-functionalized derivatives **2a-2c** were prepared by exhaustive alkylation of the appropriate mono-functionalized derivatives with a large excess of ethyl bromoacetate in refluxing acetone (yield: 69-82%).

To have a more versatile procedure for the synthesis of mono-penta type mixed ligands, the monophenol pentaester **3** was chosen as a key intermediate with the intent of utilizing benzyl ether moiety as a protecting group. Monobenzyl ether pentaethyl ester **2d** was prepared from monobenzyl ether of *p*-tert- butylcalix[6]arene⁶ by exhaustive alkylation with ethyl bromoacetate (K₂CO₃/acetone). Subsequent removal of benzyl group by treatment with trimethylsilyl bromide⁵ yield monophenol-pentaester **3** (65%). We attempted to introduce the desired functional group into the remaining phenol group of the *p*-tert- butylcalix[6]arene by alkylation with a suitable reagent but to no avail partly due to the steric congestion in monophenol **3**. For all the prepared ligands, ethoxycarbomethyl substituent was chosen as penta-part because of their widely investigated and relatively well-defined ionophoric properties toward many interesting guests.⁷