

## 단 신

### Structure and Vibrational Frequencies of $\text{CH}_2=\text{NbH}_2$ , a Major Reaction Product of Methane Activation by Laser-Ablated Nb atoms

조 한 국\*  
인천대학교 화학과  
(2006. 7. 25 접수)

### Structure and Vibrational Frequencies of $\text{CH}_2=\text{NbH}_2$ , a Major Reaction Product of Methane Activation by Laser-Ablated Nb atoms

Han-Gook Cho\*  
Department of Chemistry, University of Incheon, 177 Dohwa-dong, Nam-ku, Incheon, 402-749, South Korea  
(Received July 25, 2006)

주제어: 메틸리덴 착물, 아고스틱 상호작용, 레이저 애블레이션, 니오비움, 매트릭스

**Keywords:** Methylidene Complex, Agostic Interaction, Laser Ablation, Nb, Matrix

Viable activation of methane is a long standing goal for chemists. Recent studies show that methane activation by laser-ablated transition metal atoms leads to formation of the simple insertion product  $\text{CH}_3\text{-MH}$ ,<sup>1-7</sup> and in the case of Group 4 and 6 metals,  $\alpha\text{-H}$  transfer also gives the methylidene dihydride complex  $\text{CH}_2=\text{MH}_2$ . Electronic structure calculations have shown agostic stabilization of these methylidene complexes through  $\text{CH}_2$  distortion and unequal M-H bond lengths.<sup>3-7</sup> The activation of a second methane molecule gave new dimethylmetaldihydrides  $(\text{CH}_3)_2\text{MH}_2$  in the case of Group 4, but Group 6 allowed a second  $\alpha\text{-H}$  transfer to form the methylidyne  $\text{CH}\equiv\text{MH}_3$ . Such simple methylidene and methylidyne complexes provide a model for structures and substituent effects for the much larger high-oxidation state complexes previously synthesized.<sup>8</sup>

Niobium alkylidene complexes have been mostly synthesized by  $\alpha\text{-H}$  abstraction reactions; however, attempts to provide a Nb methylidyne complex have not been successful giving related complexes instead.<sup>8</sup> Recently reaction of laser-ablated niobium atoms with methane has been carried out and  $\text{CH}_3\text{-NbH}$ ,

$\text{CH}_2=\text{NbH}_2$ ,  $\text{CH}\equiv\text{NbH}_3^-$ , and  $(\text{CH}_3)_2\text{NbH}_2$  are identified in the matrix infrared spectra.<sup>9</sup> Questions, however, still remain in the structure of the methylidene hydride complex,  $\text{CH}_2=\text{NbH}_2$ . Spectroscopic evidences apparently show that the two Nb-H bonds are apparently equal, which are compared with the highly asymmetric  $\text{MH}_2$  groups of Group 4 and 6 metal methylidenes.<sup>3-7</sup> We report here an investigation with electronic structure computations for the structure of the smallest possible niobium methylidene hydride complex ( $\text{CH}_2=\text{NbH}_2$ ).

The frequencies of the product absorptions observed from reaction of Nb with methane are listed in Table 1.<sup>9</sup> The most distinctive absorptions are  $\text{NbH}_2$  stretching bands, which are strong and appear in a relatively clean area. Deuteration leads to variations in frequency for all the absorptions, and particularly the hydrogen stretching frequencies show large isotopic shifts (H/D ratios of 1.39). On the other hand,  $^{13}\text{C}$  substitution results in almost no change in the high frequency region, but gives sizable shifts in the low frequency region.

Without Nb hydrides<sup>10</sup> in the matrix spectra, the two strong Nb-H stretching absorptions with 1:2 inten-

Table 1. Frequencies of product absorptions observed from reactions of methane with Nb in excess argon<sup>a</sup>

Group	CH <sub>4</sub>	CD <sub>4</sub>	<sup>13</sup> CH <sub>4</sub>	CH <sub>2</sub> D <sub>2</sub>
<b>i</b>	1611.4	1159.1	1611.4	1611.0
	1680.7	1206.0	1680.7	1680.7, <b>1666.4</b> , 1652.5
<b>m</b>	1652.5	1189.9	1652.5	1205.5, <b>1198.4</b> , 1188.7
	805.4	736.8	784.5	
	547.1		546.6	
	480.4		480.4	
<b>a</b>	1542.5	1103.7	1542.2	1524.3
	1489.5	1076.1	1489.5	1506.7, 1488.2
<b>di</b>	1717.5	1231.5	1717.5	1717.1, 1694.5, 1686.3
	1686.3	1213.0	1686.3	1217.8, 1212.8

<sup>a</sup>All frequencies are taken from Ref [9] and in cm<sup>-1</sup>. Stronger absorptions are bold. **i**, **m**, **a**, and **di** stand for CH<sub>3</sub>-NbH, CH<sub>2</sub>=NbH<sub>2</sub>, CH≡NbH<sub>3</sub><sup>+</sup>, and (CH<sub>3</sub>)<sub>2</sub>NbH<sub>2</sub>, respectively.

sity ratio and other **m** absorptions (in Table 1) indicate that CH<sub>2</sub>=NbH<sub>2</sub> is readily formed in reaction of Nb with CH<sub>4</sub>. Computation results also indicate that the small Schrock carbene is in fact most stable among the plausible products.<sup>9</sup> CH<sub>2</sub>D<sub>2</sub> provides further valuable diagnostic information for the product structure. Deuteration of one of the hydrogen atoms for the NbH<sub>2</sub> group effectively breaks down the coupling between the two hydrogen stretching motions.

The hydrogen stretching frequencies of CHD=NbHD in Table 1, which are at the middle of the symmetric and antisymmetric hydrogen stretching pairs, suggest that CH<sub>2</sub>=NbH<sub>2</sub> in its doublet ground state has a C<sub>s</sub> structure with two equal Nb-H bonds. Extensive calculations have been carried out at the

levels of B3LYP, BPW91, MP2, and CCSD with 6-311++G(2d,p) and 6-311++G(3df,3pd) bases.<sup>11</sup> Interestingly enough, at all levels of calculation performed in this study, two stable conformations are identified for CH<sub>2</sub>=NbH<sub>2</sub>, the C<sub>i</sub> and C<sub>s</sub> structures (Fig. 1). The T1 diagnostic values for the CCSD calculations are, however, fairly high (~0.06), suggesting that single-reference calculations are not enough for accurate determination of the structure.

Notice that the C<sub>i</sub> structure shown in Fig. 1, which is slightly more stable (less than 3 kcal/mol) than the C<sub>s</sub> structure, is agostically distorted. The agostic interaction, observed as inclination of the C-H bond toward the transition metal center, is in fact quite common among high oxidation-state complexes. Clot and Eisenstein<sup>12</sup> and Scherer and McGrady<sup>13</sup>




Fig. 1. Two stable (C<sub>i</sub> and C<sub>s</sub>) structures of CH<sub>2</sub>=NbH<sub>2</sub> in the ground doublet states. The C<sub>i</sub> and C<sub>s</sub> structure shown here are calculated both at the level of CCSD/6-311++G(3df, 3pd)/SDD (for Nb). The bond lengths and angles are in Å and degrees.

Table 2. Observed and calculated fundamental frequencies of  $\text{CH}_2=\text{NbH}_2$  isotopomers for configuration 2 ( $C_s$ ) in the Ground  $^2A'$  Electronic State<sup>a</sup>

Approximate Description	$\text{CH}_2=\text{NbH}_2$							$\text{CD}_2=\text{NbD}_2$			$^{13}\text{CH}_2=\text{NbH}_2$			$\text{CHD}=\text{NbHD}$		
	Freq.						Int.	Freq.			Int.	Freq.			Int.	Int.
	Obs.	B3LYP anhar <sup>b</sup>	B3LYP	CCSD	BPW91	MP2		Obs.	CCSD	Int.		Obs.	CCSD	Int.	Obs.	CCSD
$\nu_1$ A' C-H str.		3097.8	3225.4	3272.1	3162.6	3305.2	7		2425.1	10		3260.7	6		2556.2	18
$\nu_2$ A' C-H str.		2473.5	2599.6	2529.7	2459.6	2517.6	21		1842.4	8		2523.8	21		2394.9	11
$\nu_3$ A' Nb-H str.	1680.7	1716.9	1771.9	1787.7	1749.9	1941.6	289	1206.0	1269.0	146	1680.7	1787.7	289	1666.4	1780.6	373
$\nu_4$ A' $\text{CH}_2$ scis.		1336.8	1362.5	1448.2	1350.8	1520.1	36		1100.6	15		1441.8	37		1350.5	26
$\nu_5$ A' C=Nb str.	805.4	828.4	853.8	889.8	852.5	1531.3	88	736.8	802.0	66	784.5	866.3	86	805.4	849.2	71
$\nu_6$ A' $\text{NbH}_2$ scis.		606.8	654.6	696.3	676.3	796.4	72		515.8	24		694.8	73		594.9	67
$\nu_7$ A' $\text{NbH}_2$ wag		511.9	550.7	643.1	585.7	744.6	23		473.4	29		639.6	20		565.1	16
$\nu_8$ A' $\text{CH}_2$ rock	547.1	420.5	505.7	531.1	546.6	536.7	77		381.9	37	546.6	530.7	77		476.7	66
$\nu_9$ A'' Nb-H str.	1652.5	1701.8	1761.7	1774.3	1744.2	1879.5	470	1189.9	1267.1	243	1652.5	1774.3	469	1198.4	1268.4	197
$\nu_{10}$ A'' $\text{CH}_2$ wag		602.5	629.3	684.4	622.0	794.5	45		537.0	22		678.5	45		547.7	20
$\nu_{11}$ A'' $\text{NbH}_2$ twist	480.4	404.3	457.2	458.7	469.2	549.8	66		330.7	36	480.4	457.8	65		357.1	39
$\nu_{12}$ A'' $\text{CH}_2$ twist		143.4	309.9	291.8	310.8	473.3	0		207.7	0		291.8	0		267.7	1

<sup>a</sup>Calculated with 6-311++G(3df, 3pd)/SDD (for Nb). Frequencies and intensities are in  $\text{cm}^{-1}$  and  $\text{km/mol}$ . Infrared intensities are calculated values with CCSD. <sup>b</sup>Anharmonic frequencies calculated with B3LYP.

recently reported that agostic stabilization has little or no dependence on C-H-M electron donation but arises from delocalization of the C=M bonding electrons. The  $C_1$  structure varies considerably depending on the level of theory. B3LYP density functional underestimates the agostic distortion, while MP2 method and BPW91 density functional overestimate it. More rigorous CCSD method gives an intermediate result.

On the other hand, the  $C_s$  structure is quite unique with two equal Nb-H bonds. Unlike the  $C_1$  structure, the  $C_s$  structure hardly varies regardless of the level of theory, which coincides with the experimental evidences for the two equal Nb-H bonds. It is interesting that the  $CH_2$  group is also highly distorted (*Fig. 1*), in fact even more than in the  $C_1$  structure. The calculated harmonic and anharmonic frequencies at various level of theory for the  $C_s$  structure are listed in *Table 2*. Among them, the BPW91 frequencies of 1749.9, 1744.2, 852.5, 546.6, and 469.2  $cm^{-1}$  best match with the observed values of 1680.7, 1652.5, 805, 547, 480.4  $cm^{-1}$ . The frequencies of the other methane isotopomers are also compared with the CCSD frequencies in *Table 2*, showing reasonable agreements.

The calculated vibrational characteristics, while showing reasonable agreements, also show systematic discrepancies with the experimental values. For example, the calculated symmetric and anti-symmetric Nb-H stretching frequencies are mostly too close to each other in comparison with the observed values (1680.7 and 1652.5  $cm^{-1}$ , 28.2  $cm^{-1}$  apart). However, unlike other methods, the MP2 frequencies are too far apart (62.1  $cm^{-1}$ ). The level of agreement between the experimental and calculated values varies markedly in the low frequency region with the calculation methods. Evidently, more accurate molecular structure of  $CH_2=NbH_2$  ought to be determined while the  $C_s$  structure is better in line than the  $C_1$  structure with the experimental results.<sup>9</sup>

We have shown that B3LYP, BPW91, MP2, and CCSD calculations with reasonably large bases give two possible configurations ( $C_1$  and  $C_s$ ) with similar energies for  $CH_2=NbH_2$ . While the  $C_1$  configuration is consistent with those of previously studied Group

4 and 6 methyldene complexes, the  $C_s$  structure with two equal Nb-H bonds reasonably reproduces the observed frequencies and particularly the observed isotopic variations. However, the calculated vibrational characteristics also show systematic discrepancies with the observed characteristics, and they, along with the high T1 diagnostic value and the illusive two configurations, request further theoretical examinations such as multi-reference computations.

**Acknowledgment.** We gratefully acknowledge financial support from University of Incheon.

## REFERENCES

1. Billups, W. E.; Konarski, M. M.; Hauge, R. H.; Margrave, J. L. *J. Am. Chem. Soc.* **1980**, *102*, 7394.
2. Wang, G.; Chen, M.; Zhou, M. *Chem. Phys. Letts.* **2005**, *412*, 46.
3. Cho, H.-G.; Wang, X.; Andrews, L. *J. Am. Chem. Soc.* **2005**, *127*, 465.
4. Cho, H.-G.; Wang, X.; Andrews, L. *Organometallics* **2005**, *24*, 2854.
5. Andrews, L.; Cho, H.-G.; Wang, X. *Inorg. Chem.* **2005**, *44*, 4834.
6. Cho, H.-G.; Andrews, L. *J. Am. Chem. Soc.* **2005**, *127*, 8226.
7. Cho, H.-G.; Andrews, L.; Marsden, C. *Inorg. Chem.* **2005**, *44*, 7634.
8. Schrock, R. R. *Chem. Rev.* **2002**, *102*, 145.
9. Cho H.-G.; Andrews, L. *J. Phys. Chem. A* **2006**, *110*, 3886.
10. (a) Van Zee, R. J.; Weltner, W., Jr. *J. Chem. Phys.* **1995**, *102*, 4367. (b) Wang, X.; Andrews, L. unpublished data (Nb, Ta+H<sub>2</sub>).
11. Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu,

- G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. *Gaussian 03*, Revision B.04, Gaussian, Inc.: Pittsburgh, PA, 2003 and references therein.
12. Clot, E.; Eisenstein, O. Agostic Interactions from a Computational Perspective. In *Structure and Bonding, Computational Inorganic Chemistry*; Kaltzoyannis, N., McGrady, E., Eds.; Springer-Verlag: Heidelberg, Germany, 2004; pp 1-36.
13. Scherer, W.; McGrady, G. S. *Angew. Chem., Int. Ed.* **2004**, *43*, 1782.
-