

A New Amaryllidaceae Alkaloid from the Bulbs of *Lycoris radiata*Ji Young Lee,^{†,‡} Mi-Ran Cha,[†] Ji Eun Lee,^{†,‡} Jinhee Kim,[†] Heeyeong Cho,[†] Woo Kyu Park,[†]
Sang Un Choi,[†] and Shi Yong Ryu^{†,‡,*}[†]Korea Research Institute of Chemical Technology, Daejeon 305-600, Korea. *E-mail: syryu@kRICT.re.kr[‡]Graduate School of New Drug Discovery and Development, Chungnam National University, Daejeon 305-764, Korea

Received July 30, 2014, Accepted August 29, 2014

Key Words : *Lycoris radiata*, Amaryllidaceae alkaloids, Acetylcholinesterase inhibition

Lycoris radiata (red spider lily) is a bulbous perennial plant in Amaryllidaceae widely distributed in Korea, Japan and China. It is commonly used as an ornamental flower or sometimes as an antidote in Traditional Chinese Medicine (TCM).¹ The species is also known as a botanical source of galantamine, a clinically used medicine for the treatment of Alzheimer's diseases and various other memory impairments.² Actually, the bulbs of the species contains a lot of isoquinoline-based amine components, so called Amaryllidaceae alkaloids, which have been reported to possess a variety of pharmacological activities such as anti-cancer,³ anti-viral,⁴ anti-acetylcholinesterase⁵ and anti-inflammatory⁶ activities. Amaryllidaceae alkaloids are produced almost by plants of galanthus genus⁷ in Amaryllidaceae and are classified with three distinct scaffolds, lycorine (3-9, 14), haemanthamine (1-2, 10, 15-18) and galanthamine (11-13) series, which are classified depending on the pattern of

oxidative phenolic coupling. For the purpose of searching for bioactive alkaloids from natural resources, extensive phytochemical investigation of the bulbs extract of *L. radiata* had undertaken and finally resulted in the isolation of a new amaryllidaceae alkaloid (1), together with seventeen related alkaloids (2-18). The chemical structures of isolated 1-18 (Fig. 1) were established by spectroscopic analyses as 6-hydroxytazettine (1), tazettine (2), lycorine (3), 2-*O*-acetyllycorine (4), homolycorine (5), 9-*O*-demethylhomolycorine (6), hippeastrine (7), *O*-methyllycorine (8), 2- α -hydroxy-6-*O*-methyloduline (9), haemanthidine (10), galantamine (11), dihydrogalantamine (12), lycoramine N-oxide (13), lycosinineB (14), ismine (15), trisphaeridine (16), 3-epi-macronine (17), and 6-*O*-methylpretazettine (18),⁸⁻¹⁸ respectively. In the present paper, we describe briefly the isolation and identification of a new Amaryllidaceae alkaloid (1), as well as the inhibitory activities of isolated alkaloids (1-18)


**Figure 1.** Structures of isolated compounds 1-18.

Table 1. ^1H and ^{13}C NMR Spectroscopic Data of **1** and **2**^a

Position	1 (δ_{H})	2 (δ_{H})	1 (δ_{C})	2 (δ_{C})
1	5.64 (1H, d, $J = 10.4$ Hz)	5.64 (1H, d, $J = 10.4$ Hz)	127.6	128.7
2	6.00 (1H, d, $J = 10.4$ Hz)	6.16 (1H, d, $J = 10.4$ Hz)	129.5	130.6
3	4.06 (1H, m)	4.16 (1H, ddd, $J = 10.4, 5.8, 1.8$ Hz)	72.6	72.9
4	2.23 (1H, m)	2.24 (1H, dddd, $J = 13.5, 5.8, 4.0, 1.1$ Hz)	25.7	26.7
	1.68 (1H, m)	1.65 (1H, ddd, $J = 13.5, 10.4, 2.3$ Hz)		
4a	3.01 (1H, m)	2.88 (1H, m)	67.4	70.0
6	4.41 (1H, s)	3.32 (1H, d, $J = 10.6$ Hz)	93.0	65.6
		2.70 (1H, d, $J = 10.6$ Hz)		
6a			101.4	102.1
8α	5.03 (1H, $J = 14.5$ Hz)	4.98 (1H, d, $J = 14.7$ Hz)	62.1	62.1
8β	4.63 (1H, $J = 14.5$ Hz)	4.65 (1H, d, $J = 14.7$ Hz)		
8a			125.6	125.5
9	6.48 (1H, s)	6.52 (1H, s)	103.8	103.9
10			146.5	146.4
11			146.6	146.6
12	6.91 (1H, s)	6.88 (1H, s)	109.6	109.4
12a			127.2	128.0
12b			47.9	49.9
-OCH ₂ O-	5.92 (2H, d, $J = 3.7$ Hz)	5.92 (2H, s)	100.9	100.9
-OCH ₃	3.45 (3H, s)	3.48 (3H, s)	56.1	56.1
-NCH ₃	2.62 (3H, s)	2.42 (3H, s)	38.8	41.9

^aAssignments are based on DEPT, HSQC, and HMBC experiments, and chemical shifts are given in ppm

on acetylcholinesterase, *in vitro* assay.

Compound **1** was obtained as a white amorphous powder. The molecular formula of **1** was established as C₁₈H₂₁NO₆, m/z 347.1424 (calcd. 347.1369) by HREIMS. The ^1H -NMR data, particularly signals in lower field were quite similar with those of tazettine (**2**), which suggested that **1** is a kind of congener of **2** (Table 1). **2** has been isolated previously from other plants in Amaryllidaceae family, *i.e.*, *Narcissus tazetta* and *Pancratium maritimum*, and the chemical structure of **2** as well as the absolute configuration was established by X-ray diffraction analysis.¹⁹

The ^1H -NMR data of **1** implied the presence of a tetra-substituted benzene ring (δ_{H} 6.91, 6.48), a pair of olefin (δ_{H} 6.00, 5.64), a methylenedioxy group (δ_{H} 5.92), and another methylene group (δ_{H} 5.03, 4.63). However, when the ^1H -NMR spectrum of **1** was compared with that of **2**, a pair of methylene signals of H-6 observed at δ_{H} 3.32 and 2.70 in ^1H -NMR of **2** was disappeared, instead, a singlet proton signal at δ_{H} 4.41 was observed in ^1H -NMR of **1**. These ^1H -NMR results suggested that one of methylene protons of **2** was replaced with hydroxyl group to yield **1**, and this assumption was supported by the mass differences ($\Delta 16$) between **1** and **2**.

Thus, All proton and carbon signals of **1** were completely assigned by the aid of two-dimensional NMR experiments such as COSY, DEPT, HSQC, HMBC, and NOESY, which led to the conclusion that one of methylene protons at C-6 of **2** was replaced with hydroxyl group to become **1**, thus **1** is established as 6-hydroxytazettine. In HMBC experiment of **1**, the H-6 (δ_{H} 4.41) signal was observed to give correlations with C-12b, C-6a, and *N*-methyl carbon, respectively (Fig.




Figure 2. Key HMBC ($\text{H} \rightarrow \text{C}$) (a) and NOESY ($\text{H} \leftrightarrow \text{H}$) (b) correlations of **1**.

2). Supposing that the stereochemistry of C-4a, C-12b and C-6a of **2** were identical with those of **1**, the relative configuration of **1**, particularly the C-6 chirality was proposed by the NOESY experiment; the signal of H-6 was correlated with H-4a and H-8 α , which strongly suggested a β -orientation of the hydroxyl group attached at C-6 (Fig. 2) and the *R* configuration of C-6. Thus, **1** was proposed as 6 β -hydroxytazettine as illustrated in Figure 1 and Figure 2. The other purified components (**2**-**18**) were identified by direct comparison of their spectral data with those in the literatures.⁸⁻¹⁸ All isolated Amaryllidaceae alkaloids (**1**-**18**) were evaluated for the inhibitory effect on acetylcholinesterase (EC 3.1.1.7. electric eel) using acetylcholine as substrate *in vitro*²⁰ and it was observed that galantamine (**11**) and ismine (**15**) exhibited a significant inhibitory effect on the acetylcholinesterase with IC₅₀ values of 27.9 and 109.7 μM , when other isolated compounds showed poor inhibitions (IC₅₀ > 300 μM) on acetylcholinesterase, *in vitro* assay.

Experimental Section

General Experimental Procedures. NMR spectra were obtained by a Bruker AM 300 and 500 spectrometers using TMS as an internal standard for ^1H NMR, ^{13}C NMR, DEPT, COSY, HSQC, HMBC, and NOESY. Preparative-HPLC was performed on a Futecs P-4000 system with a Shim-pack prep-ODS(H) kit column (5 μm , 20 mm $\times$ 25 cm). Isolation and purification was also carried out using a medium-pressure liquid chromatographic (MPLC) system [BUCHI pump Module C-601, silica gel 60 (230-400 mesh, Merck), ODS (Cosmosil 140 C₁₈)].

Plant Material. The bulbs of *L. radiata* were collected on November 2013 at the flower garden of Korea Research Institute of Chemical Technology (KRICT) and were authenticated by Dr. Young sup Kim. A voucher specimen (KR1315) was deposited at the herbarium of the KRICT.

Extraction and Isolation. The dried bulbs (6 Kg) of *L. radiata* were soaked twice in 90 L of methanol at room temperature for 7 days. The concentrated methanol extract (700 g) was suspended in 4 L of distilled water and adjusted to pH 12.0 with 1 N NaOH, and then extracted successively with equal volume of ethylacetate (EtOAc) and *n*-butanol (BuOH). The resultant EtOAc layer and BuOH layer was collected and extracted with equal volume 5% citric acid, respectively. The 5% citric acid layer was pooled up and neutralized with 1 N NaOH to pH 12.0 and then extracted with BuOH, which gave 39 g of the alkaloids rich fraction. The alkaloids rich fraction (39 g) was subjected to column chromatography on silica gel eluted with dichloromethane : MeOH (0.5 to 100% MeOH gradient) to yield four fractions (Fr. 1 - Fr. 4). Fr. 3 (6.2 g) was dissolved in methanol and allowed to crystallize at room temperature to give compound **3** (2.33 g). Fr. 1 (1.6 g) was subjected to column chromatography with NH₂ gel (Merck Lichroprep NH₂) eluted with a hexane : EtOAc (0.5 to 100% EtOAc gradient) to give seven fractions (Fr. 11 - Fr. 17). Fr. 12 afforded compound **18** (2.3 mg) by repeated preparative HPLC (50 to 100% MeOH). Fr. 13 was further purified through sephadex LH-20 chromatography and preparative HPLC (30 to 90% MeOH) to yield compounds **17** (4.5 mg), **1** (2.2 mg), **16** (3.4 mg). Fr. 14 was chromatographed on a sephadex LH-20 columns to give compounds **4** (5.1 mg) and **15** (7.5 mg). Fr. 2 (5.5 g) was further purified to column chromatography on silica gel eluted with dichloromethane : MeOH (1 to 100% MeOH gradient) to give four fractions (Fr. 21 - Fr. 24). Fr. 22 was purified further by preparative HPLC (50-90% MeOH) to yield compound **5** (38.7 mg) and **14** (3.1 mg). Compound **2** (27.4 mg) was crystallized from Fr. 24 in MeOH. Fraction 3 (6.17 g) was isolated to five fractions (Fr. 31 - Fr. 35). Fr. 33 was repeatedly chromatographed by preparative HPLC (30 - 90% MeOH) to afford compounds **8** (125.0 mg), **9** (5.0 mg), **11** (14.1 mg) and **7** (44.2 mg). Fr. 35 was subjected to column chromatography on silica gel eluted with dichloromethane : MeOH (1 to 20% MeOH gradient) to yield compound **6**

(150.0 mg). Fr. 4 was subjected to column chromatography with silica gel eluted with dichloromethane : MeOH (2 to 50% MeOH gradient) to give five fractions (Fr. 41 - Fr. 45). Fr. 44 was isolated by preparative HPLC (10 to 80% MeOH) and finally purified through sephadex LH-20 to afford compounds **10** (34.2 mg), **12** (74.0 mg) and **13** (4.8 mg).

6-Hydroxytazettine (1): White amorphous powder; $[\alpha]_D^{20} +35.9$ (c 0.05, CHCl₃); UV (MeOH) λ_{max} 240, 291 nm; ^1H NMR (CDCl₃, 500 MHz); ^{13}C NMR (CDCl₃, 125 MHz) (Table 1); HREIMS m/z 347.1424 (calcd. for C₁₈H₂₁NO₆, 347.1369).

Acknowledgments. This research was supported by the basic research project (KK-1403-C0) of KRICT and also by a grant of the Technology Innovation Program (10038744) of Korea Evaluation Institute of Industrial Technology (KEIT) funded by Ministry of Trade, Industry & Energy, Republic of Korea.

References

- Huang, S. D.; Zhang, Y.; He, H. P.; Li, S. F.; Tang, G. H.; Chen, D. Z.; Cao, M. M.; Di, Y. T.; Hao, W. J. *Chinese Journal of Natural Medicines* **2013**, *11*, 406.
- Berkov, S.; Georgieva, L.; Kondakova, V.; Atanassov, A.; Viladomat, F.; Bastida, J.; Codina, C. *Biotechnol. & Biotechnol. Eq.* **2009**, *23*, 1170.
- Antonio, E.; Alexander, K. *Phytochem. Rev.* **2009**, *8*, 449.
- He, J.; Qi, W. B.; Wang, L.; Tian, J.; Jiao, P. R.; Liu, G. Q.; Ye, W. C.; Liao, M. *Influenza Other Respir. Viruses* **2013**, *6*, 922.
- Susana, L.; Jaume, B.; Francesc, V.; Carles, C. *Life Sciences* **2002**, *71*, 2521.
- Saltan, C. G.; Dahadir, A. O.; Sever, Y. B.; Ozbek, H. *Fitoterapia* **2012**, *83*, 81.
- Nehir, U. *Phytochem. Rev.* **2007**, *6*, 125.
- Pham, L. H.; Grundemann, E.; Wagner, J.; Bartoszek, M.; Dopke, W. *Phytochemistry* **1999**, *51*, 327.
- Evidente, A.; Cicala, M. R.; Giudicianni, I.; Randazzo, G.; Riccio, R. *Phytochemistry* **1983**, *22*, 581.
- Szlavik, L.; Gyuris, A.; Minarovits, J.; Forgo, P.; Molnar, J.; Hohmann, J. *Planta Med.* **2004**, *70*, 871.
- Kihara, M.; Konishi, K.; Xu, L.; Kobayashi, S. *Chem. Pharm. Bull.* **1991**, *39*, 1849.
- Almanza, Z. R.; Fernandez, J. M.; Wakori, E. W. T.; Viladomat, F.; Codina, C.; Bastida, J. *Phytochemistry* **1996**, *43*, 1375.
- Jegorov, A.; Buchta, M.; sedmera, P.; Kuzma, M.; Havlicek, V. *J. Mass Spectrom.* **2006**, *41*, 544.
- Kobayashi, S.; Satoh, K.; Numata, A.; Shingu, T.; Kihara, M. *Phytochemistry* **1991**, *30*, 675.
- Yang, Y.; Huang, S. X.; Zhao, Y. M.; Zhao, Q. S.; Sun, H. D. *Helvetica Chimica Acta* **2005**, *88*, 2550.
- Kihara, M.; Koike, Y.; Imakura, Y.; Kida, K.; Shingu, T.; Kobayashi, S. *Chem. Pharm. Bull.* **1987**, *35*, 1070.
- Vdovin, A. D.; Kadyrov, K. H.; Yagudaev, M. R.; Allayarov, K. B.; Nistryan, A. K. *Chemistry of Natural Compounds* **1981**, *17*, 279.
- Cabezas, F.; Ramirez, A.; Viladomat, F.; Codina, C.; Bastida, J. *Chem. Pharm. Bull.* **2003**, *51*, 315.
- Ide, S.; Sener, B.; Temizer, H.; Konukol, S. *Cryst. Res. Technol.* **1996**, *31*, 617.
- Lee, J. Y.; Cha, M. R.; Choi, C. W.; Kim, Y. S.; Lee, B. H.; Ryu, S. Y. *Kor. J. Pharmacogn.* **2012**, *43*, 122.