

L-Ascorbic Acid로부터 2-Deoxy-L-Ribose의 효과적인 합성

金寬洙* · 安榮熙 · 許恩榮 · 李宜宰

연세대학교 이과대학 화학과

(1994. 8. 22 접수)

Efficient Synthesis of 2-Deoxy-L-Ribose Starting from L-Ascorbic Acid

Kwan Soo Kim*, Yeong Hee Ahn, Eun Young Hurh, and Eui Jae Lee


Department of Chemistry, Yonsei University, Seoul 120-749, Korea

(Received August 22, 1994)

The chirality of the sugar-backbone units in natural nucleic acids is responsible for the formation of the high-order structure of the nucleic acids and for their functions as well. Organisms on the earth utilize only the D-sugar. Nevertheless, calculations¹ and other studies² suggest that modified nucleic acids based on L-sugar recognize complementary nucleic acids. Furthermore, oligonucleotides composed of 2-deoxy-L-ribose (2-deoxy-L-*erythro*-pentose **1**) show resistance to digestion by certain nucleases²⁻⁴. Enantio-DNA (DNA having 2-deoxy-L-ribose) and meso-DNA (DNA having an alternating sequence of L-sugar and D-sugar) are, therefore, valuable tools for studying protein-DNA interactions and are promising antisense agents⁵⁻⁷. In this regard, there still remains a need for the efficient synthetic method for 2-deoxy-L-ribose. Among a few known methods for the synthesis of **1**⁸⁻¹⁰, only the glycal method developed by Deriaz *et al.*⁹ has been used in practice. Herein we report a new efficient method for the synthesis of compound **1** starting from L-ascorbic acid (**2**).

L-Ascorbic acid (**2**) was converted to 5,6-O-isopropylidene derivative **3** in 95% yield by treatment with acetyl chloride in acetone¹¹. Oxidation of **3** with hydrogen peroxide in the presence of calcium carbonate¹² afforded the threonic acid derivative **4**¹³ in 72% yield. Compound **4** was transformed

into the methyl ester **5**¹³ in 95% yield with methyl iodide and sodium bicarbonate in dimethylacetamide. The secondary hydroxyl group of **5** was tosylated with tosyl chloride and triethylamine in methylene chloride to give compound **6**¹⁴ in 90% yield. The reduction of tosylate **6** with sodium borohydride in methanol and subsequent epoxidation of the resulting 1,2-hydroxytosylate with potassium carbonate were carried out in one pot to afford epoxide **7**¹⁵ in 63% yield. Reaction of compound **7** with lithiated 1,3-dithiane in THF at -40 °C provided white solid **8**¹⁶ in 70% yield. To a solution of thioacetal **8** (0.18 g, 0.68 mmol) in water (16 ml)-acetone (16 ml) was added 1.0 N HCl (1.0 ml) at room temperature. After stirring for 30 min, HgO (0.66 g, 3.04 mmol) and HgCl₂ (0.80 g, 2.96 mmol) were added to the reaction mixture and the stirring continued for further 2 h at 40 °C. The reaction mixture was filtered and acetone was removed *in vacuo*. To the remaining aqueous solution was added Na₂S (0.86 g, 3.56 mmol) and precipitated HgS was removed. The volume of aqueous solution was reduced to a half by freeze-drying. Isopropyl alcohol (20 ml) was added to the aqueous solution and precipitated sodium chloride was removed by filtration. The filtrate was evaporated to give a pale yellow syrup which was crystallized in vacuum after one week. Recrystallization of the



crude crystal from ethyl acetate gave pure 2-deoxy-L-ribose (**1**, 0.06 g, 70%)¹⁷. The overall yield of the present procedure is 18%. The present method is superior over Deriaz's glycol method⁹ at least in two aspects: the overall yield and the cost of the starting material.

We thank Yonsei University for financial support. Support from OCRC is also acknowledged.

REFERENCES

1. Ashley, G. W. *J. Am. Chem. Soc.* **1992**, *114*, 9731.
2. Fujimori, S.; Shudo, K.; Hashimoto, Y. *J. Am. Chem. Soc.* **1990**, *112*, 7436.
3. Damha, M. J.; Giannaris, P. A.; Marfey, P.; Reid, L. S. *Tetrahedron Lett.* **1991**, *32*, 2573.
4. Asseline, U.; Hau, J.-F.; Czernecki, S.; Diguarher, T. L.; Perlat, M.-C.; Valery, J.-M.; Thuong, N. T. *Nucleic Acids Res.* **1991**, *19*, 4067.
5. Goodchild, J. *Bioconjugate Chem.* **1990**, *1*, 16.
6. Uhlmann, E.; Peyman, A. *Chem. Rev.* **1990**, *90*, 543.
7. Beaucage, S. L.; Iyer, R. P. *Tetrahedron* **1993**, *49*, 6123.
8. Meisenheimer, J.; Jung, H. *Ber.* **1927**, *60*, 1462.
9. Deriaz, R. E.; Overend, W. G.; Stacey, M.; Teece, E. G.; Wiggins, L. F. *J. Chem. Soc.* **1949**, 1879.
10. Schimmel, S. D.; Beville, R. D. *Annal. Biochem.* **1970**, *37*, 385.
11. Jackson, K.; Jones, J. *Can. J. Chem.* **1969**, *42*, 2498.
12. Isbell, H. S.; Frush, H. L. *Carbohydr. Res.* **1979**, *72*, 301.
13. Wei, C. C.; Bjernardo, S. De.; Tengi, J. P.; Borgese, J.; Weigle, M. *J. Org. Chem.* **1985**, *50*, 3462.
14. Compound **6**: mp. 56~58 °C; ¹H NMR (80 MHz, CDCl₃) δ 1.22 (s, 6H), 2.37 (s, 3H), 3.63 (s, 3H), 3.88~3.92 (m, 2H), 4.28~4.48 (m, 1H), 4.75~4.81 (d, 1H), 7.19~7.81 (2d, 4H).
15. Le Merr, Y.; Gravier-Pelletier, C.; Dumas, J.; Depéay, J. C. *Tetrahedron Lett.* **1990**, *31*, 1003.
16. Compound **8**: mp. 71 °C; ¹H NMR (300 MHz, CDCl₃) δ 1.17 (s, 3H), 1.36 (s, 3H), 1.17~2.10 (m, 4H), 2.21 (s, 1H), 2.76~2.84 (m, 5H), 3.85~4.03 (m, 3H), 4.19~4.24 (m, 1H).
17. Compound **1**: mp. 89 °C (lit.⁹ 90 °C); [α]_D +55.0° (c 0.27, H₂O) (lit.⁹ +58°); ¹H NMR (200 MHz, D₂O) δ 1.64~2.50 (m, 2H), 3.57~4.42 (m, 4H), 5.30~5.65 (m, 1H).